

Group algebras and semigroup algebras defined by permutation relations of fixed length

Ferran Cedó*

Eric Jespers†

Georg Klein

Abstract

Let H be a subgroup of Sym_n , the symmetric group of degree n . For a fixed integer $l \geq 2$, the group G presented with generators $x_1, x_2, \dots, x_n$ and with relations $x_{i_1}x_{i_2} \cdots x_{i_l} = x_{\sigma(i_1)}x_{\sigma(i_2)} \cdots x_{\sigma(i_l)}$, where σ runs through H , is considered. It is shown that G has a free subgroup of finite index. For a field K , properties of the algebra $K[G]$ are derived. In particular, the Jacobson radical $\mathcal{J}(K[G])$ is always nilpotent, and in many cases the algebra $K[G]$ is semiprimitive. Results on the growth and the Gelfand-Kirillov dimension of $K[G]$ are given. Further properties of the semigroup S and the semigroup algebra $K[S]$ with the same presentation are obtained, in case S is cancellative. The Jacobson radical is nilpotent in this case as well, and sufficient conditions for the algebra to be semiprimitive are given.

Keywords: group ring, semigroup ring, finitely presented, group, semigroup, Jacobson radical, semiprimitive, primitive.

Mathematics Subject Classification: Primary 16S15, 16S34, 16S36, 20M05; Secondary 20M25, 16N20.

1 Introduction

Motivated by a general interest in finitely presented algebras with homogeneous defining relations, in [1], the authors initiated the study of the monoid algebras $K[S_{n,l}(H)]$, where $l \geq 2$ is an integer and H is a subgroup of Sym_n and $S_{n,l}(H)$ is the monoid presented with generators $x_1, x_2, \dots, x_n$ and with relations $x_{i_1}x_{i_2} \cdots x_{i_l} = x_{\sigma(i_1)}x_{\sigma(i_2)} \cdots x_{\sigma(i_l)}$, for all $i_1, i_2, \dots, i_l \in \{1, 2, \dots, n\}$ and all $\sigma \in H$. In the present article, we obtain further results on algebras of this type. Let $G_{n,l}(H)$ be the universal group of $S_{n,l}(H)$, that is the group presented with the “same” generators and relations. Recall that a subgroup H of Sym_n is semi-regular if the stabilizer of i in H is trivial for all $i \in \{1, 2, \dots, n\}$. In [1, Theorems 2.3 and 2.5] it is shown that $S_{n,l}(H)$ is cancellative if and only if H is semi-regular and abelian, and in this case $S_{n,l}(H)$ embeds in $G_{n,l}(H)$. For simplicity, we will denote $S_{n,l}(H)$ by S , and $G_{n,l}(H)$ by G .

In Section 2, we prove that, for arbitrary H , the group G has a free subgroup of finite index. A similar result for S was obtained in [1]. In Section 3, we show that the Jacobson radical, $\mathcal{J}(K[G])$, is nilpotent. We also prove that if H is transitive, then $K[G]$ is a Noetherian PI-algebra, and the Gelfand-Kirillov dimension and the classical Krull dimension are both equal to 1. In Section 4, we show that if H is semi-regular and abelian, then $\mathcal{J}(K[S])$ is nilpotent of index at most $|G/M|^2$,

*Research partially supported by a grant of MICIIN (Spain) MTM2011-28992-C02-01.

†Research supported in part by Onderzoeksraad of Vrije Universiteit Brussel and Fonds voor Wetenschappelijk Onderzoek (Belgium).

where M is a normal subgroup of finite index of G . This is a generalization of [1, Theorem 3.1], in that the assumption that H is transitive is removed.

2 Groups $G_{n,l}(H)$ defined by permutation relations

For a subgroup H of Sym_n and an integer $l \geq 2$, let $S_{n,l}(H) = \langle x_1, x_2, \dots, x_n \mid x_{i_1}x_{i_2} \cdots x_{i_l} = x_{\sigma(i_1)}x_{\sigma(i_2)} \cdots x_{\sigma(i_l)}, \sigma \in H, i_1, \dots, i_l \in \{1, \dots, n\} \rangle$. We consider the universal group $G_{n,l}(H)$ with the same presentation as $S_{n,l}(H)$. In [1], it was shown that if H is not abelian, then $S_{n,l}(H)$ is not embedded in $G_{n,l}(H)$.

Lemma 2.1 *In $G_{n,l}(H)$ the following equalities hold for an integer $u \geq 1$:*

$$\begin{aligned} x_{j_1}x_{j_2} \cdots x_{j_u}x_{k_u}^{-1}x_{k_{u-1}}^{-1} \cdots x_{k_1}^{-1} &= x_{\sigma(j_1)}x_{\sigma(j_2)} \cdots x_{\sigma(j_u)}x_{\sigma(k_u)}^{-1}x_{\sigma(k_{u-1})}^{-1} \cdots x_{\sigma(k_1)}^{-1}, \\ x_{j_1}^{-1}x_{j_2}^{-1} \cdots x_{j_u}^{-1}x_{k_u}x_{k_{u-1}} \cdots x_{k_1} &= x_{\sigma(j_1)}^{-1}x_{\sigma(j_2)}^{-1} \cdots x_{\sigma(j_u)}^{-1}x_{\sigma(k_u)}x_{\sigma(k_{u-1})} \cdots x_{\sigma(k_1)}, \end{aligned}$$

where $j_1, j_2, \dots, j_u, k_u, k_{u-1}, \dots, k_1 \in \{1, 2, \dots, n\}$.

Proof. Let z be the smallest integer such that $zl > u$. Because of the defining relations of $S_{n,l}(H)$, in its universal group for all $\sigma \in H$ we have $x_{j_1}x_{j_2} \cdots x_{j_u}x_1^{zl-u} = x_{\sigma(j_1)}x_{\sigma(j_2)} \cdots x_{\sigma(j_u)}x_{\sigma(1)}^{zl-u}$. Hence $x_{\sigma(j_u)}^{-1}x_{\sigma(j_{u-1})}^{-1} \cdots x_{\sigma(j_1)}^{-1}x_{j_1}x_{j_2} \cdots x_{j_u} = x_{\sigma(1)}^{zl-u}x_1^{u-zl}$. Therefore

$$\begin{aligned} x_{\sigma(k_u)}^{-1}x_{\sigma(k_{u-1})}^{-1} \cdots x_{\sigma(k_1)}^{-1}x_{k_1}x_{k_2} \cdots x_{k_u} &= x_{\sigma(1)}^{zl-u}x_1^{u-zl} \\ &= x_{\sigma(j_u)}^{-1}x_{\sigma(j_{u-1})}^{-1} \cdots x_{\sigma(j_1)}^{-1}x_{j_1}x_{j_2} \cdots x_{j_u}, \end{aligned}$$

and hence

$$x_{j_1}x_{j_2} \cdots x_{j_u}x_{k_u}^{-1}x_{k_{u-1}}^{-1} \cdots x_{k_1}^{-1} = x_{\sigma(j_1)}x_{\sigma(j_2)} \cdots x_{\sigma(j_u)}x_{\sigma(k_u)}^{-1}x_{\sigma(k_{u-1})}^{-1} \cdots x_{\sigma(k_1)}^{-1}.$$

The other equality can be proved similarly. ■

Lemma 2.2 *In $G_{n,l}(H)$ the following equalities hold:*

$$\begin{aligned} x_{\sigma_1(1)}x_{\sigma_2(1)} \cdots x_{\sigma_{l-1}(1)}x_{\sigma_l(j)} &= x_1^{l-1}x_jx_1^{1-l}x_{\sigma_1^{-1}\sigma_2(1)}x_{\sigma_1^{-1}\sigma_3(1)} \cdots x_{\sigma_1^{-1}\sigma_l(1)}, \\ x_{\sigma_1(1)}x_{\sigma_2(1)} \cdots x_{\sigma_{l-1}(1)}x_{\sigma_l(j)}^{-1} &= x_1^{l-1}x_j^{-1}x_1^{1-l}x_{\sigma_l\sigma_{l-1}^{-1}(1)}x_{\sigma_l\sigma_{l-1}^{-1}\sigma_1(1)} \cdots x_{\sigma_l\sigma_{l-1}^{-1}\sigma_{l-2}(1)}, \end{aligned}$$

where $j \in \{1, 2, \dots, n\}$ and $\sigma_1, \dots, \sigma_l \in H$.

Proof. By the defining relations of $G_{n,l}(H)$ and Lemma 2.1, we have

$$\begin{aligned}
& x_{\sigma_1(1)}x_{\sigma_2(1)} \cdots x_{\sigma_{l-1}(1)}x_{\sigma_l(j)} \\
&= x_1x_{\sigma_1^{-1}\sigma_2(1)} \cdots x_{\sigma_1^{-1}\sigma_{l-1}(1)}x_{\sigma_1^{-1}\sigma_l(j)}x_1^{1-l}x_1^{l-1} \\
&= x_1^2x_{\sigma_2^{-1}\sigma_3(1)} \cdots x_{\sigma_2^{-1}\sigma_{l-1}(1)}x_{\sigma_2^{-1}\sigma_l(j)}x_{\sigma_2^{-1}\sigma_1(1)}^{1-l}x_1^{l-1} \\
&= x_1^3x_{\sigma_3^{-1}\sigma_4(1)} \cdots x_{\sigma_3^{-1}\sigma_{l-1}(1)}x_{\sigma_3^{-1}\sigma_l(j)}x_{\sigma_3^{-1}\sigma_1(1)}^{2-l}x_{\sigma_2^{-1}\sigma_1(1)}^{1-l}x_1^{l-1} \\
&\vdots \\
&= x_1^{l-1}x_{\sigma_{l-1}^{-1}\sigma_l(j)}x_{\sigma_{l-1}^{-1}\sigma_1(1)}^{-2}x_{\sigma_{l-2}^{-1}\sigma_1(1)}^{-1} \cdots x_{\sigma_2^{-1}\sigma_1(1)}^{-1}x_1^{l-1} \\
&= x_1^{l-1}x_jx_{\sigma_l^{-1}\sigma_1(1)}^{-1}x_{\sigma_{l-1}^{-1}\sigma_1(1)}^{-1} \cdots x_{\sigma_2^{-1}\sigma_1(1)}^{-1}x_1^{l-1} \\
&= x_1^{l-1}x_jx_1^{-1}x_{\sigma_1^{-1}\sigma_l\sigma_{l-1}^{-1}\sigma_1(1)}^{-1} \cdots x_{\sigma_1^{-1}\sigma_l\sigma_2^{-1}\sigma_1(1)}^{-1}x_{\sigma_1^{-1}\sigma_l(1)}^{l-1} \\
&= x_1^{l-1}x_jx_1^{-2}x_{\sigma_1^{-1}\sigma_{l-1}\sigma_{l-2}^{-1}\sigma_1(1)}^{-1} \cdots x_{\sigma_1^{-1}\sigma_{l-1}\sigma_2^{-1}\sigma_1(1)}^{-1}x_{\sigma_1^{-1}\sigma_{l-1}(1)}^{l-2}x_{\sigma_1^{-1}\sigma_l(1)} \\
&\vdots \\
&= x_1^{l-1}x_jx_1^{1-l}x_{\sigma_1^{-1}\sigma_2(1)}x_{\sigma_1^{-1}\sigma_3(1)} \cdots x_{\sigma_1^{-1}\sigma_l(1)},
\end{aligned}$$

and the first equality is proved. Now applying the first equality for $\sigma_l\sigma_{l-1}^{-1}, \sigma_l\sigma_{l-1}^{-1}\sigma_1, \dots, \sigma_l\sigma_{l-1}^{-1}\sigma_{l-2}, \sigma_l \in H$, we obtain

$$x_{\sigma_l\sigma_{l-1}^{-1}(1)}x_{\sigma_l\sigma_{l-1}^{-1}\sigma_1(1)} \cdots x_{\sigma_l\sigma_{l-1}^{-1}\sigma_{l-2}(1)}x_{\sigma_l(j)} = x_1^{l-1}x_jx_1^{1-l}x_{\sigma_1(1)}x_{\sigma_2(1)} \cdots x_{\sigma_{l-1}(1)},$$

and thus the second equality follows. $\blacksquare$

Theorem 2.3 Assume that the different orbits under the action of H are $X_{k_1}, X_{k_2}, \dots, X_{k_r}$, where X_k denotes the orbit of k , and $k_1 = 1$. Then the subgroup F of $G_{n,l}(H)$ generated by $x_1, x_{k_2}, \dots, x_{k_r}$ is free of rank r , and for every $g \in G_{n,l}(H)$, there exist $w \in F$ and $\sigma_1, \sigma_2, \dots, \sigma_{l-1} \in H$ such that

$$g = wx_{\sigma_1(1)}x_{\sigma_2(1)} \cdots x_{\sigma_{l-1}(1)}.$$

Proof. Let $G = G_{n,l}(H)$. Let G_1 be the normal closure of $\text{gr}(x_{k_j}x_{\sigma(k_j)}^{-1} \mid k_j \in \{1, k_2, \dots, k_r\}, \sigma \in H)$ in G . We consider $\bar{G} = G/G_1$. Then $\bar{G} = \text{gr}(x_1, x_2, \dots, x_n \mid x_{i_1}x_{i_2} \cdots x_{i_l} = x_{\sigma(i_1)}x_{\sigma(i_2)} \cdots x_{\sigma(i_l)}, x_{k_j} = x_{\sigma(k_j)}, \sigma \in H, i_1, \dots, i_l \in \{1, \dots, n\}, k_j \in \{1, k_2, \dots, k_r\})$. The second relation in the presentation of $\bar{G}$ implies the first, and $\bar{G} = \text{gr}(X_{k_1}, X_{k_2}, \dots, X_{k_r})$ is a free group of rank r . Consider the natural epimorphism $\pi : G \rightarrow \bar{G}$. If there was a non-trivial relation in G , of type $w = 1$ with w a reduced word in $\{x_1^{\pm 1}, x_{k_2}^{\pm 1}, \dots, x_{k_r}^{\pm 1}\}$, then we would have $\pi(w) = 1$, a contradiction because $\bar{G}$ is a free group. Thus the first part of the result follows.

For the second part, we will use induction on the length of the reduced words in $\{x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_n^{\pm 1}\}$ that represent an element $g \in G$. For length 0 we have $1 = x_1^{1-l}x_1^{l-1}$ and for length 1, according to Lemma 2.1 and the defining relations, we have

$$\begin{aligned}
x_{\sigma(k)} &= x_{\sigma(k)}x_1^{-1}x_1^{2-l}x_1^{l-2}x_1 = x_kx_{\sigma^{-1}(1)}^{-1}x_1^{2-l}x_1^{l-2}x_1 \\
&= x_kx_1^{-1}x_{\sigma(1)}^{2-l}x_{\sigma(1)}^{l-2}x_{\sigma(1)} = x_kx_1^{-1}x_1^{2-l}x_1^{l-2}x_{\sigma(1)}, \\
x_{\sigma(k)}^{-1} &= x_{\sigma(k)}^{-1}x_1^{1-l}x_1^{l-1} = x_k^{-1}x_{\sigma^{-1}(1)}^{1-l}x_1^{l-1} = x_k^{-1}x_1^{1-l}x_{\sigma(1)}^{l-1}.
\end{aligned}$$

Let $s \geq 1$ and assume the result is true for all $g \in G$ such that the length of some reduced word in $\{x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_n^{\pm 1}\}$ that represents g is s . Let

$$x_{\sigma_1(j_1)}^{\epsilon_1}x_{\sigma_2(j_2)}^{\epsilon_2} \cdots x_{\sigma_{s+1}(j_{s+1})}^{\epsilon_{s+1}},$$

be a reduced word with $\epsilon_u \in \{1, -1\}$, $j_u \in \{1, k_2, \dots, k_r\}$ and $\sigma_u \in H$, for $1 \leq u \leq s+1$. By induction hypothesis there exist $w \in F$ and $\tau_1, \dots, \tau_{l-1} \in H$ such that

$$x_{\sigma_1(j_1)}^{\epsilon_1} x_{\sigma_2(j_2)}^{\epsilon_2} \cdots x_{\sigma_{s+1}(j_{s+1})}^{\epsilon_{s+1}} = w x_{\tau_1(1)} \cdots x_{\tau_{l-1}(1)} x_{\sigma_{s+1}(j_{s+1})}^{\epsilon_{s+1}}.$$

Now by Lemma 2.2, the result follows. ■

3 The algebra $K[G_{n,l}(H)]$

In this section we study the algebraic structure of the group algebra $K[G_{n,l}(H)]$ over a field K , for any subgroup H of Sym_n . We denote $G_{n,l}(H)$ by G .

Theorem 3.1 *Let F be the free subgroup of G defined in the statement of Theorem 2.3. Then $[G : F] \leq |H|^{l-1}$ and $(\mathcal{J}(K[G]))^{[G:F]} = 0$. Furthermore, if the characteristic of K does not divide $[G : F]$, then $\mathcal{J}(K[G]) = 0$.*

Proof. By Theorem 2.3, we know that $[G : F] \leq |H|^{l-1}$. Then by a result of Villamayor [7, Theorem 7.2.7], $(\mathcal{J}(K[G]))^{[G:F]} \subseteq (\mathcal{J}(K[F])) \cdot K[G]$. Because F is a free group, $\mathcal{J}(K[F]) = 0$, and hence $(\mathcal{J}(K[G]))^{[G:F]} = 0$. If the characteristic of K does not divide $[G : F]$, then by the same result [7, Theorem 7.2.7], we have $\mathcal{J}(K[G]) = (\mathcal{J}(K[F])) \cdot K[G] = 0$. ■

Theorem 3.2 (a) *If H is transitive, then $K[G]$ is a Noetherian PI-algebra and*

$$\text{GKdim}(K[G]) = 1 = \text{clKdim}(K[G]).$$

(b) *If H is not transitive, then $K[G]$ has exponential growth.*

Proof.

- (a) If H is transitive, by Theorem 2.3, $K[G]$ is a finitely generated module over $K[\text{gr}(x_1)] = K[x_1, x_1^{-1}]$. Thus $K[G]$ is a Noetherian PI-algebra and $\text{GKdim}(K[G]) = 1$. By [5, Theorem 23.4], $\text{clKdim}(K[G]) = \text{GKdim}(K[G]) = 1$.
- (b) If H is not transitive, by Theorem 2.3, G contains a free subgroup of rank 2. Thus $K[G]$ has exponential growth.

■

4 The algebra $K[S_{n,l}(H)]$

In this section, we obtain a generalization of [1, Theorem 3.1].

To do so, we recall Lemma 2.2 and Theorem 2.5 of [1]. The first part is the monoid version of Theorem 2.3.

Proposition 4.1 *Let H be a subgroup of Sym_n . Then $S = S_{n,l}(H)$ contains a free submonoid FM , with basis $\{x_1, x_{k_2}, \dots, x_{k_r}\}$ (one element from each orbit on the generating set), and a finite subset T such that $S = \bigcup_{t \in T} \text{FM}t = \bigcup_{t \in T} t\text{FM}$. If, furthermore, H is abelian and semi-regular, then S is a submonoid of its universal group $G = G_{n,l}(H)$.*

Recall that a semigroup P is said to be right reversible if $Pa \cap Pb \neq \emptyset$ for all $a, b \in P$.

We need the following elementary technical lemma.

Lemma 4.2 *A right reversible submonoid of a free monoid is contained in a cyclic group.*

Proof. Let B be a right reversible submonoid of a free monoid FM. Let $|x|$ denote the length of x in a chosen basis X of the free monoid FM. Let $s, t \in B$. Then $Bs \cap Bt \neq \emptyset$. Thus $as = bt$ for some $a, b \in B$. Because FM is free, it follows that if $|s| \geq |t|$, then $s = vt$ for some $v \in \text{FM}$. In particular, if $x, y \in B$ and $|x| = |y|$, then $x = y$.

Clearly, there exist positive integers n and m such that $|s^n| = |t^m|$. Since $s^n, t^m \in B$, it follows that $s^n = t^m$.

Let FG denote the free group with basis X . Consider the group $C = \langle s, s^{-1}, t, t^{-1} \rangle \subseteq \text{FG}$. Because a subgroup of a free group is free, we get that C is a free group of rank at most two. As $s^n = t^m$, the rank has to be one and thus C is cyclic. So, any two elements of B commute and thus B generates an abelian subgroup in the free group in FG. So, this group is cyclic. ■

Because of Theorem 2.3, $G = G_{n,l}(H)$ has a free subgroup of finite index. Let M be a normal subgroup of G that is of finite index and a free group.

Theorem 4.3 *Let H be an abelian and semi-regular subgroup of Sym_n . Let $G = G_{n,l}(H)$ be the universal group of $S = S_{n,l}(H)$. Then, $\mathcal{J}(K[S])$ is nilpotent of index at most $|G/M|^2$. If $\text{char}(K) = 0$ or $\text{char}(K) = p$ and $p \nmid |G/M|$, then $\mathcal{J}(K[S]) = 0$ and $\mathcal{J}(K[G]) = 0$.*

Proof. Obviously, one can consider $K[G]$ in a natural way as a G/M graded ring. Its component of degree e (the identity of G/M) is $K[M]$. By Proposition 4.1, $K[S]$ is a homogeneous subring of $K[G]$ and hence it also is a G/M -graded ring, with component of degree e the subring $K[M \cap S]$. As a subsemigroup of M , the semigroup $M \cap S$ is a two-unique product monoid and thus $\mathcal{J}(K[S \cap M]) = 0$. Since $K[S]$ is graded by a finite group, it then follows that $\mathcal{J}(K[S])^{m^2} = 0$, where $m = |G/M|$ (this follows from [3, 4] and Lemma 20 in [2]). This proves the first part of the theorem.

To prove the second part, assume $\text{char}(K) = 0$ or $\text{char}(K) = p$ and $p \nmid m$. We prove the result by contradiction. So, assume $\mathcal{J}(K[S]) \neq 0$. Then, by [6, Corollary 1], there exists a left and right reversible submonoid P of S such that $\mathcal{J}(K[P]) \neq 0$.

We claim that $P \cap M$ also is a right reversible monoid. Indeed, let $s, t \in P \cap M$. Then $Ps \cap Pt \neq \emptyset$ and thus $p_1s = p_2t$, for some $p_1, p_2 \in P$. Hence, $(p_2^{m-1}p_1)s = (p_2^m)t$. Clearly, $p_2^m \in P \cap M$ and thus $p_2^{m-1}p_1 \in M \cap P$. Thus, $(P \cap M)s \cap (P \cap M)t \neq \emptyset$. So, indeed, $P \cap M$ is right reversible.

By Lemma 4.2, $P \cap M$ is contained in a cyclic subgroup of M . Let $Q = PP^{-1}$ be the group of fractions of P . Then $Q \cap M$ is normal in Q and $Q \cap M$ is a cyclic group. Let $p \in P$. Clearly, p acts by conjugation on $Q \cap M$. Since the latter is cyclic, this conjugation either is the trivial map or it maps an element to its inverse. Consequently, if $f \in P \cap M$ and $p \in P$, then $pf = f^{-1}p$ or $pf = fp$. By a degree argument, the former is impossible. So $pf = fp$. Hence, $P \cap M$ is central in P and thus $Q \cap M = (P \cap M)(P \cap M)^{-1}$ is a central subgroup of Q of finite index. Actually the index is a divisor of m .

Hence Q is a finite conjugacy group, with an infinite cyclic central subgroup of finite index that is a divisor of m . By [7, Lemma 4.1.4] such a group has a finite normal subgroup, say N , such that Q/N is cyclic. Clearly, N is isomorphic to a subgroup of G/M and thus $|N|$ is a divisor of m . Hence, by [7, Theorem 7.3.1], $\mathcal{J}(K[Q]) \subseteq \mathcal{J}(K[N])K[Q]$. Moreover, by assumption, either

$\text{char}(K) = 0$ or $p \nmid m$. So, by Maschke's Theorem, $\mathcal{J}(K[Q]) = 0$. Further, by [5, Corollary 11.5], $\mathcal{J}(K[P]) = \mathcal{J}(K[Q]) \cap K[P]$. So $\mathcal{J}(K[P]) = 0$, a contradiction. ■

References

- [1] F. Cedó, E. Jespers and G. Klein, Finitely presented monoids and algebras defined by permutation relations of abelian type, II, J. Pure App. Algebra 219 (2015), 1095–1102.
- [2] M. V. Clase and E. Jespers, On the Jacobson radical of semigroup graded rings. J. Algebra 169 (1994), no. 1, 79–97.
- [3] M. Cohen and S. Montgomery, Group-graded rings, smash products, and group actions. Trans. Amer. Math. Soc. 282 (1984), no. 1, 237–258.
- [4] M. Cohen and S. Montgomery, Addendum to: "Group-graded rings, smash products, and group actions" [Trans. Amer. Math. Soc. 282 (1984), no. 1, 237–258; MR0728711 (85i:16002)]. Trans. Amer. Math. Soc. 300 (1987), no. 2, 810–811.
- [5] J. Okniński, Semigroup Algebras, Marcel Dekker, 1991.
- [6] J. Okniński, Algebras of cancellative semigroups, Bull. Austral. Math. Soc. 49 (1994), 165–170.
- [7] D.S. Passman, The Algebraic Structure of Group Rings, Wiley, New York, 1977.

F. Cedó

Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193 Bellaterra (Barcelona), Spain
cedo@mat.uab.cat

E. Jespers

Department of Mathematics
Vrije Universiteit Brussel
Pleinlaan 2, 1050 Brussel, Belgium
efjesper@vub.ac.be

G. Klein

Department of Mathematics
Vrije Universiteit Brussel
Pleinlaan 2, 1050 Brussel, Belgium
gklein@vub.ac.be